

On the one-dimensional representations of the general linear supergroup

I.M. Trishin

Abstract. Because of its multiplicativity, the Berezinian is the character of the one-dimensional representation of the general linear supergroup. We give an explicit construction of this representation on a space of tensors. Similarly, we construct the representation such that its character is the inverse of the Berezinian.

0 Introduction

The following result belongs to the basic ones in the classic case. Let V be an n -dimensional linear space over a field K of zero characteristic. Then the space $\text{Alt}(\underbrace{V \otimes \cdots \otimes V}_n)$ of the skew-symmetric tensors of degree n is the one-dimensional module over the general linear group $GL V$. The character of such representation of $GL V$ is the determinant. In the present paper we construct a pair of analogous representations on spaces of tensors for the supercase. The character of one of these representations is the Berezinian. The other character is the inverse of the Berezinian.

Outline the way to our construction in general. Let G be a Grassman algebra with a countable set of generators, $V = V_0 \oplus V_1$ a free $\mathbb{Z}_2$ -graded G -module of dimension $m|n$, where $m = \dim_G V_0$, $n = \dim_G V_1$. Suppose W_{λ_h} , W_{λ_g} are the irreducible $GL V$ -modules defined by the partitions $\lambda_h = (\underbrace{n+1, \dots, n+1}_m)$, $\lambda_g = (\underbrace{n, \dots, n}_{m+1})$ (respectively) and b is a formal element such that for an arbitrary $\mathcal{A} \in GL V$ we have $\mathcal{A}(b) = \text{Ber } \mathcal{A} \cdot b$, where A is a matrix of $\mathcal{A}$. We show that the $GL V$ -modules W_{λ_h} and $b \cdot W_{\lambda_g}$ are isomorphic (see Theorem 2.1). To prove this theorem we use a natural correspondence between the sets of the base vectors of W_{λ_h} and W_{λ_g} . The tensor $\tilde{b}$ generating

the one-dimensional GLV -module is constructed of the element b and these base vectors of W_{λ_h} and W_{λ_g} (see Theorem 3.1). By analogy we construct the one-dimensional GLV -module such that its character is the inverse of the Berezinian.

This paper is organized as follows. We introduce the main concepts in §1. In §2 we prove that $W_{\lambda_h} \simeq b \cdot W_{\lambda_g}$ and obtain an important corollary of this result (see Theorem 2.2). Then in §3 the one-dimensional representations of GLV are constructed.

In relation to the present paper, the result of H.M. Khudaverdian and Th.Th. Voronov must be mentioned. In [1], the Berezinian is expressed as the ratio of the two Hankel determinants such that these are the characters of the representations of GLV defined by the partitions λ_h and λ_g . On one hand, this is a simple corollary of Theorem 2.1, on the other directly implies this theorem (unfortunately, the last easy observation have come too late to be useful for the author).

The author wishes to thank Th.Th. Voronov for useful discussions.

1 The basic concepts and auxiliary results

Let K be a field of characteristic zero, $G = G_0 \oplus G_1$ the Grassman algebra over K with a countable set of generators.

By definition, the notion of G -module includes the property to be free.

Let $V = V_0 \oplus V_1$ be a finite-dimensional $\mathbb{Z}_2$ -graded G -bimodule and the structures of the left and right G -modules on V are compatible (see, for example, [2], [3]). Denote $m = \dim_G V_0$, $n = \dim_G V_1$. The pair $m|n$ is the *dimension* of V .

We suppose that any operation of changing base of V is even.

Let $\text{End } V$ be the algebra of all G -linear mappings from V to V . The algebra $\text{End } V$ is isomorphic to the algebra $M = M_0 \oplus M_1$ of block matrices with Grassman elements. The even component M_0 of M is called the *full matrix superalgebra* and is denoted by $M_{m,n}$. Actually the elements of $M_{m,n}$ are all matrices of the form $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$, where A_{11} , A_{22} are square G_0 -matrices of orders m and n respectively and A_{12} , A_{21} are rectangular G_1 -matrices of corresponding orders.

Let GLV and $GL_{m,n}$ be the groups of all invertible elements of $\text{End } V$ and $M_{m,n}$ respectively. We have the isomorphism $GLV \simeq GL_{m,n}$. The group

$GL_{m,n}$ (GLV) is the *general linear supergroup*.

Recall that for any $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \in GL_{m,n}$ the *Berezinian* of A is given by the formula

$$\begin{aligned} \text{Ber } A &= \det(A_{11} - A_{12}A_{22}^{-1}A_{21}) \det A_{22}^{-1} = \\ &= \det A_{11} \det(A_{22} - A_{21}A_{11}^{-1}A_{12})^{-1} \end{aligned}$$

(see [4]). The main feature of the Berezinian is its multiplicativity.

Denote

$$T_l(V) = \underbrace{V \otimes \cdots \otimes V}_l,$$

where $l = 1, 2, \dots$. Clearly, $T_l(V)$ is a G -module. By definition, for any $\mathcal{A} \in GLV$, $v_i \in V$, $l = 1, 2, \dots$ we have

$$\mathcal{A}(v_1 \cdots v_l) = \mathcal{A}(v_1) \cdots \mathcal{A}(v_l). \quad (1.1)$$

Action (1.1) of GLV on decomposable tensors is extended to the whole space $T_l(V)$ by linearity and we see that $T_l(V)$ is a GLV -module.

In what follows we consider the action induced by GLV on some submodules of $T_l(V)$. Nevertheless again the corresponding operator is denoted by $\mathcal{A}$. This cannot confuse us forasmuch as from context is clear, what of submodules is considered.

Let S_l be the symmetric group on the elements $1, 2, \dots, l$.

By definition, for an arbitrary $\sigma \in S_l$, $v_i \in V$, $i = 1, 2, \dots, l$ put

$$\sigma(v_1 \cdots v_l) = v_{\sigma^{-1}(1)} \cdots v_{\sigma^{-1}(l)}. \quad (1.2)$$

Then $T_l(V)$ is a left S_l -module.

To determine irreducible GLV -submodules of $T_l(V)$ the special element of the group algebra $K[S_l]$, the so-called *Young symmetrizer*, can be used. Recall this notion.

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be positive integers such that $\lambda_1 \geq \dots \geq \lambda_k$ and $l = \lambda_1 + \dots + \lambda_k$. Then $\lambda = (\lambda_1, \dots, \lambda_k)$ is the *partition* of l . The corresponding *Young diagram* λ has λ_i boxes in its i -th row. The *Young tableau* T_λ is obtained from λ by filling its boxes with the numbers $1, 2, \dots, l$ moving by the rows from left to right and from top to bottom. For example, if $\lambda = (4, 2, 1)$, then T_λ has the form

1	2	3	4
5	6		
7			

Let $C(T_\lambda)$ ($R(T_\lambda)$) be the subgroup of S_l that preserves the numbers of T_λ within their columns (rows, respectively). Then

$$e_T = \sum_{\substack{\sigma \in C(T), \\ \tau \in R(T)}} \text{sgn}(\sigma)\tau\sigma$$

is the *Young symmetrizer* and $W_\lambda = e_{T_\lambda} T_l(V)$ is the irreducible GLV -module.

Denote

$$\lambda_h = (\underbrace{n+1, \dots, n+1}_m),$$

$$\lambda_g = (\underbrace{n, \dots, n}_{m+1}).$$

In other words, λ_h is the $m \times (n+1)$ rectangle and λ_g is the $(m+1) \times n$ rectangle. In the sequel, the GLV -modules W_{λ_h} , W_{λ_g} play an important role.

Remark. One can say that W_{λ_g} , W_{λ_h} are particular cases of the so-called *external forms* (see [2]).

Denote

$$\lambda_{SR} = (\underbrace{n, \dots, n}_m),$$

$$\lambda_{LR} = (\underbrace{n+1, \dots, n+1}_{m+1}),$$

$$\lambda_{AR} = (\underbrace{n+1, \dots, n+1}_m, n).$$

According to terminology of Issaia L. Kantor the diagrams λ_{SR} , λ_{LR} , λ_{AR} are called the *small rectangle*, the *large rectangle*, the *almost rectangle* (respectively).

Evidently, to obtain λ_h (λ_g) one must add the column of height m (the row of length n) to the small rectangle. As mentioned above, in either case we obtain a kind of rectangle.

An arbitrary tensor of the form

$$e_T v_1 \cdots v_l, \quad (1.3)$$

where $v_1, v_2, \dots, v_l \in V$, has the following properties:

i) if $i_1, i_2 \in \{1, 2, \dots, l\}$ place in the same column of T then tensor (1.3) is skew-symmetric by the elements v_{i_1}, v_{i_2} , that is,

$$e_T v_1 \cdots v_l = -e_T(i_1, i_2) v_1 \cdots v_l; \quad (1.4)$$

ii) let $i_1, \dots, i_{k-1}$ be numbers that fill a column of length $(k-1)$ of T , i_k a number that belongs to a column of length not more than $(k-1)$; then the *Jacobi identity* holds, that is,

$$e_T v_1 \cdots v_l = \sum_{j=1}^{k-1} e_T(i_j, i_k) v_1 \cdots v_l \quad (1.5)$$

(see [5]). Clearly, (1.4) and (1.5) are the applications of the corresponding properties of the symmetrizer e_T .

Suppose f is a tensor of the form (1.3). We say that (i, j) is an *f-box* if this box belongs to the tableau T defining the tensor f . By analogy the concepts of an *f-row* and an *f-column* are introduced.

Let $\{e_1, \dots, e_m\}, \{\varepsilon_1, \dots, \varepsilon_n\}$ be some bases of the G -modules V_0 and V_1 respectively. Denote

$$\Omega_0 = \{e_1, \dots, e_m\}, \quad \Omega_1 = \{\varepsilon_1, \dots, \varepsilon_n\}, \quad \Omega = \Omega_0 \cup \Omega_1.$$

We say that an element $f \in T_l(V)$ is an Ω -*tensor* if f has the form (1.3), where $v_i \in \Omega$ for $i = 1, 2, \dots, l$.

Recall that we use the unique way of filling of Young diagrams (see above). Thus there is a one-to-one correspondence between Ω -tensors and fillings of the corresponding Young diagrams with the vectors $v_i \in \Omega$.

A filling of a rectangular Young diagram by vectors $v_i \in \Omega$ is *canonical* if every even basis vector $e_i \in \Omega_0$ cannot be placed anywhere except boxes of the i -th row and every odd basis vector $\varepsilon_j \in \Omega_1$ cannot be placed anywhere except boxes of the j -th column. In other words, a filling of a rectangular

diagram is canonical if a box (i, j) that is within the small rectangle can be filled by e_i or by ε_j only.

For example, if the dimension of V is $2|1$, $\Omega_0 = \{e_1, e_2\}$, $\Omega_1 = \{\varepsilon\}$, then the canonical fillings of the small rectangle are the following:

$$\begin{array}{|c|} \hline e_1 \\ \hline e_2 \\ \hline \end{array}, \quad \begin{array}{|c|} \hline e_1 \\ \hline \varepsilon \\ \hline \end{array}, \quad \begin{array}{|c|} \hline \varepsilon \\ \hline e_2 \\ \hline \end{array}, \quad \begin{array}{|c|} \hline \varepsilon \\ \hline \varepsilon \\ \hline \end{array}.$$

Suppose λ is a rectangular Young diagram filled with elements of Ω in a canonical way. Then the contents of boxes that do not belong to the small rectangle is defined uniquely. More precisely, if these boxes belong to the i -th row, where $i \in \{1, 2, \dots, m\}$ (j -th column, where $j \in \{1, 2, \dots, n\}$), then they are filled with e_i (with ε_j respectively).

In particular, the canonical fillings of the diagrams λ_g , λ_h have the following properties:

- i) the $(n + 1)$ -th column of the diagram λ_h is filled with the vectors $e_1, \dots, e_m$ from the top down;
- ii) the $(m + 1)$ -th row of the diagram λ_g is filled with the vectors $\varepsilon_1, \dots, \varepsilon_n$ from left to right.

For example, if the dimension of V is $1|2$, $\Omega_0 = \{e\}$, $\Omega_1 = \{\varepsilon_1, \varepsilon_2\}$, then the canonical fillings of the diagram λ_g are the following:

$$\begin{array}{|c|c|} \hline e & e \\ \hline \varepsilon_1 & \varepsilon_2 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline \varepsilon_1 & e \\ \hline \varepsilon_1 & \varepsilon_2 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline e & \varepsilon_2 \\ \hline \varepsilon_1 & \varepsilon_2 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline \varepsilon_1 & \varepsilon_2 \\ \hline \varepsilon_1 & \varepsilon_2 \\ \hline \end{array}.$$

Evidently, the concept of canonical filling is defined for diagrams that do not contain the large rectangle. From the other hand, if a diagram contains the large rectangle, then tensor (1.3) is equal to zero (see, for example, [2]).

Remark. The concept of canonical filling is introduced in [2] and can be applied not only to rectangular diagrams.

We say that an Ω -tensor f is *canonical* if the filling of the corresponding diagram is canonical.

By Λ_g , Λ_h denote the sets of all canonical tensors for the diagrams λ_g , λ_h . Note that there is one-to-one correspondence between the sets Λ_g , Λ_h : the corresponding elements have the same filling within the small rectangle. Elements of the sets Λ_g , Λ_h are denoted by g_i and h_j respectively, where

$i, j = 1, 2, \dots, k_\Lambda$, $k_\Lambda = 2^{mn}$. Moreover we suppose that elements of Λ_g , Λ_h with the same numbers are corresponding, that is, for every $i \in \{1, 2, \dots, k_\Lambda\}$ tensors g_i , h_i have the same filling of the small rectangle.

Theorem 1.1. *The sets Λ_g , Λ_h are the bases of the G -modules W_{λ_g} , W_{λ_h} .*

The proof of this theorem is found in [2].

2 The isomorphism of W_{λ_h} and $b \cdot W_{\lambda_g}$ as the GLV -modules

Let b be a formal element such that for any $\mathcal{A} \in GLV$ we have

$$\mathcal{A}(b) = \text{Ber } A \cdot b, \quad (2.1)$$

where A is the matrix of $\mathcal{A}$ in some base of V . Recall that to change a base of V we use even operators only (see §1). Then taking into account the multiplicativity of Ber , one can see that the concept of the element b is well defined.

Clearly, we assume the element b is homogeneous. At the same time now we do not define if this element even or odd.

Because of the multiplicativity of Ber formula (2.1) defines the one-dimensional representation of the group GLV on the G -module generated by the element b .

Let f be an Ω -tensor. By $\kappa(f)$ denote the cardinality of the automorphism group of f -tableau by odd elements. Clearly, if an f -column does not contain any odd elements, then the corresponding factor in $\kappa(f)$ is equal to 1.

By $\rho(f)$ denote the number of Ω_1 -elements belonging to the small rectangle of f . Let α be the map from Λ_h to the rational numbers such that

$$\alpha(h_i) = \frac{(-1)^{n\rho(h_i)} \kappa(h_i)}{\kappa(g_i)}. \quad (2.2)$$

By definition, put

$$h'_i = \frac{h_i}{\alpha(h_i)}, \quad (2.3)$$

where $i = 1, 2, \dots, k_\Lambda$.

The following result is the heart of the present paper.

Theorem 2.1. *The mapping*

$$\varphi : h'_i \mapsto b \cdot g_i \quad (2.4)$$

determines the isomorphism of the $GL\ V$ -modules W_{λ_h} and $b \cdot W_{\lambda_g}$, that is, for any $\mathcal{A} \in GL\ V$ the following equality holds

$$\mathcal{A}\varphi = \varphi\mathcal{A}. \quad (2.5)$$

In the sequel we assume that $A \in GL_{m,n}$ is the matrix of $\mathcal{A} \in GL\ V$ in the base $e_1, \dots, e_m, \varepsilon_1, \dots, \varepsilon_n$.

The following proposition reduces the proof of Theorem 2.1 to some easier particular cases of the general situation.

Proposition 2.1. *To prove Theorem 2.1 it suffices to check (2.5) for the cases when the matrix A of $\mathcal{A} \in GL\ V$ belongs to one of the following classes of matrices:*

$$E_{m+n} + e_{ij}\eta_{ij}, \quad (2.6)$$

where E_{m+n} is the unit matrix of order $m+n$, e_{ij} is the matrix unit, $\eta_{ij} \in G_1$, $m+1 \leq i \leq m+n$, $1 \leq j \leq m$;

$$E_{m+n} + e_{ij}\xi_{ij}, \quad (2.7)$$

where $\xi_{ij} \in G_1$, $1 \leq i \leq m$, $m+1 \leq j \leq m+n$;

$$\text{diag}(x_1, \dots, x_m, y_1, \dots, y_n), \quad (2.8)$$

where $x_i, y_j \in G_0$;

$$E_{m+n} + e_{ij}x_{ij}, \quad (2.9)$$

where $x_{ij} \in G_0$, $1 \leq i, j \leq m$, $i \neq j$;

$$E_{m+n} + e_{ij}y_{ij}, \quad (2.10)$$

where $y_{ij} \in G_0$, $m+1 \leq i, j \leq m+n$, $i \neq j$.

The proof of this proposition is based on the following auxiliary result.

Lemma 2.1. *Any matrix $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \in GL_{m,n}$ can be represented in the form*

$$A = \begin{pmatrix} E_m & 0 \\ B_{21} & E_n \end{pmatrix} \begin{pmatrix} B_{11} & 0 \\ 0 & B_{22} \end{pmatrix} \begin{pmatrix} E_m & B_{12} \\ 0 & E_n \end{pmatrix}, \quad (2.11)$$

where $B_{11} = A_{11}$, $B_{12} = A_{11}^{-1}A_{12}$, $B_{21} = A_{21}A_{11}^{-1}$, $B_{22} = A_{22} - A_{21}A_{11}^{-1}A_{12}$, or in the form

$$A = \begin{pmatrix} E_m & C_{12} \\ 0 & E_n \end{pmatrix} \begin{pmatrix} C_{11} & 0 \\ 0 & C_{22} \end{pmatrix} \begin{pmatrix} E_m & 0 \\ C_{21} & E_n \end{pmatrix}, \quad (2.12)$$

where $C_{22} = A_{22}$, $C_{12} = A_{12}A_{22}^{-1}$, $C_{21} = A_{22}^{-1}A_{21}$, $C_{11} = A_{11} - A_{12}A_{22}^{-1}A_{21}$.

PROOF. Since $A \in GL_{m,n}$, we see that A_{11}^{-1} , A_{22}^{-1} exist (see [4]). Equalities (2.11), (2.12) can be checked directly.

This completes the proof.

PROOF OF PROPOSITION 2.1. With the isomorphism $GL V \simeq GL_{m,n}$ a composition of linear operators corresponds to the composition of matrices (clearly, the inverse statement is correct). Hence it follows from Lemma 2.1 that it suffices to check (2.5) for $\mathcal{A} \in GL V$ such that the matrix A of $\mathcal{A}$ has either the form

$$\begin{pmatrix} E_m & A_{12} \\ 0 & E_n \end{pmatrix}, \quad (2.13)$$

or

$$\begin{pmatrix} E_m & 0 \\ A_{21} & E_n \end{pmatrix}, \quad (2.14)$$

or

$$\begin{pmatrix} A_{11} & 0 \\ 0 & A_{22} \end{pmatrix}, \quad (2.15)$$

where (as above) A_{11} , A_{22} are square G_0 -matrices of orders m and n respectively, A_{12} , A_{21} are G_1 -matrices.

If i, j are integers such that $1 \leq i \leq m$, $m+1 \leq j \leq m+n$, then $e_{ij}^2 = 0$. Consequently an arbitrary matrix (2.13) can be represented as a product of matrices (2.7). Similarly any matrix (2.14) can be represented as a product of matrices (2.6).

By $\mathcal{D}$ denote the group generated by all matrices of the form (2.8) - (2.10). Also by $\bar{\mathcal{D}}$ denote the group of all matrices of the form (2.15). Just as for the regular subgroup in the classic case the set $\mathcal{D}$ is open and dense in $\bar{\mathcal{D}}$.

Thus we see that Proposition 2.1 is proved.

Applying Proposition 2.1 we use the coordinate form of equality (2.5). Let us obtain this form. With (2.1) and (2.4) we have

$$\begin{aligned}\text{Ber } A \cdot b \cdot \mathcal{A}(g_i) &= \mathcal{A}(b) \mathcal{A}(g_i) = \\ &= \mathcal{A}(bg_i) = \mathcal{A}\varphi(h'_i).\end{aligned}$$

Hence (2.5) is equivalent to

$$\varphi(\mathcal{A}(h'_i)) = \text{Ber } A \cdot b \cdot \mathcal{A}(g_i). \quad (2.16)$$

The matrix elements of the action induced by $\mathcal{A}$ on the spaces W_{λ_h} , W_{λ_g} and written in the bases $\{h'_i\}$ and $\{g_j\}$ we denote by a'_{ij} and a''_{ij} (respectively), that is,

$$\mathcal{A}(h'_j) = \sum_{i=1}^{k_\Lambda} h'_i a'_{ij}, \quad (2.17)$$

$$\mathcal{A}(g_j) = \sum_{i=1}^{k_\Lambda} g_i a''_{ij}, \quad (2.18)$$

where $1 \leq j \leq k_\Lambda$.

Using (2.17), (2.18) rewrite condition (2.16) in the form

$$\varphi\left(\sum_{i=1}^{k_\Lambda} h'_i a'_{ij}\right) = \text{Ber } A \cdot b \cdot \sum_{i=1}^{k_\Lambda} g_i a''_{ij}.$$

Hence by definition of φ we have

$$b \cdot \sum_{i=1}^{k_\Lambda} g_i a'_{ij} = \text{Ber } A \cdot b \cdot \sum_{i=1}^{k_\Lambda} g_i a''_{ij}. \quad (2.19)$$

Since the elements g_i are G -linearly independent, we see that (2.19) is equivalent to

$$a'_{ij} = \text{Ber } A \cdot a''_{ij}, \quad (2.20)$$

where $1 \leq i, j \leq k_\Lambda$. Equality (2.20) is the coordinate form of (2.5).

For an arbitrary $\mathcal{A} \in GL V$ by $A_{h'}$, A_g denote the matrices of the action induced by $\mathcal{A}$ on the spaces W_{λ_h} , W_{λ_g} and written in the bases $\{h'_i\}$, $\{g_i\}$, that is, $A_{h'} = (a'_{ij})$, $A_g = (a''_{ij})$.

Lemma 2.2. *If the matrix A of $\mathcal{A} \in GL V$ has the form (2.6), then*

$$A_{h'} = A_g.$$

For any matrix A of the form (2.6) we have $\text{Ber } A = 1$. Hence, according to Proposition 2.1, Lemma 2.2 is a part of the proof of Theorem 2.1 (see (2.20)).

PROOF OF LEMMA 2.2. By assumption, there exist integers $i \in \{1, \dots, n\}$, $j \in \{1, \dots, m\}$ such that

$$\mathcal{A}(e_j) = e_j + \varepsilon_i \eta_{ij},$$

where $\eta_{ij} \in G_1$, and $\mathcal{A}(w) = w$ for $w \in \Omega \setminus \{e_j\}$.

Let h_t, g_t be corresponding elements of the bases $\{h_i\}, \{g_i\}$, where $1 \leq t \leq k_\Lambda$, $Q_{g_t} = \{j_1, \dots, j_l\}$ (Q_{h_t}) the set of the numbers of g_t -columns (h_t -columns, respectively) that contain e_j . In other words, $r \in Q_{g_t}$ (Q_{h_t}) iff the g_t -box (h_t -box, resp.) (j, r) is filled with e_j . Evidently, $Q_{h_t} = Q_{g_t} \cup \{n+1\}$. Then we have

$$\mathcal{A}(h_t) = h_t + \sum_{r \in Q_{h_t}} \hat{h}_r \gamma_{j,r} \eta_{ij}, \quad (2.21)$$

where $\hat{h}_r$ is obtained from h_t by the substitution of ε_i for e_j placed in the h_t -box (j, r) ;

$$\gamma_{j,r} = (-1)^{n_{j,r}}, \quad (2.22)$$

where $n_{j,r}$ is the number of odd base vectors in the h_t -tableau within the interval from the box $(j, r+1)$ to the box $(m, n+1)$ (recall that we move by the tableau from left to right and from top to bottom by rows). Clearly, the tensors $\hat{h}_r$ depend on t, i, j . Nonetheless, we omit these indices in the notation to simplify the last one. In general, the tensors $\hat{h}_r$ are not canonical.

Suppose $i \neq Q_{h_t}$, that is, the h_t -box (j, i) is filled with ε_i .

Then any one of $\{\hat{h}_r | r \in Q_{h_t}\}$ is not canonical. To express $\hat{h}_r$ in terms of canonical tensors we use the Jacobi identity for the i -th column and for the box (j, r) of the $\hat{h}_r$ -tableau. Also we use (1.4) to transpose e_p and ε_r that are placed in the boxes (j, r) and (p, r) after applying (1.5).

Thus we obtain

$$\hat{h}_r = -q \hat{h}_r + \sum_{p \in P_r} h_{r,p} \beta_{r,p}, \quad (2.23)$$

where q is the number of ε_i in the i -th h_t -column (or, what is the same, in the i -th $\hat{h}_r$ -column); $P_r \subseteq \{1, \dots, m\}$ is the set of integers such that $z \in P_r$ iff e_z belongs to the i -th $\hat{h}_r$ -column and e_z does not belong to the r -th $\hat{h}_r$ -column. The tensor $h_{r,p}$ is obtained from $\hat{h}_r$ in the following two steps:

- i) the elements ε_i and e_p disposed in the $\hat{h}_r$ -boxes (j, r) and (p, i) (respectively) change places;
- ii) the elements e_p and ε_r disposed in the boxes (j, r) and (p, r) (resp.) change places (see Fig. 1).

$$\beta_{r,p} = (-1)^{\check{m}_{r,p}+1},$$

where $\check{m}_{r,p}$ is the number of pairs of odd elements that change their order under transpositions i) - ii); the additional term $(+1)$ in the exponent appears in the application of (1.4).

It is easy to see that the transformation of $\hat{h}_r$ -tableau to $h_{r,p}$ -tableau also can be realised in the following two steps: first the elements ε_i and ε_r disposed in the boxes (j, r) and (p, r) (respectively) change places; then the elements ε_i and e_p disposed in the boxes (p, r) and (p, i) (resp.) change places (see Fig. 2).




Fig. 1




Fig. 2

Then we see that

$$\beta_{r,p} = (-1)^{m_{r,p}},$$

where $m_{r,p}$ is the number of odd base vectors of the p -th row that are placed strictly between the r -th and the i -th columns.

Now it follows from (2.23) that

$$\hat{h}_r = \frac{1}{q+1} \sum_{p \in P_r} h_{r,p} \beta_{r,p}. \quad (2.24)$$

Notice that the tensors $h_{r,p}$ in (2.24) are canonical. Note also that $P_{n+1} = \emptyset$. Hence using the Jacobi identity we get $\hat{h}_{n+1} = 0$.

Combining (2.24) with (2.21) we get

$$\mathcal{A}(h_t) = h_t + \frac{1}{q+1} \sum_{r \in Q} \gamma_{j,r} \sum_{p \in P_r} h_{r,p} \beta_{r,p} \eta_{ij}, \quad (2.25)$$

where $Q = Q_{h_t} \setminus \{n+1\}$.

By assumption, $i \notin Q$. We claim that if $(r_1, p_1) \neq (r_2, p_2)$, then $h_{r_1, p_1} \neq h_{r_2, p_2}$. Indeed, if $r_1 \neq r_2$ then the sets of columns containing e_j are distinct for h_{r_1, p_1} and h_{r_2, p_2} . Further, an element e_p , where $p \in P_r$, belongs to the r -th $h_{r,z}$ -column iff $z = p$. Hence, $h_{r, p_1} \neq h_{r, p_2}$ for $p_1 \neq p_2$.

In terms of the “primed” tensors (see (2.3)) equality (2.25) takes the form

$$\mathcal{A}(h'_t) = h'_t + \frac{1}{(q+1)\alpha_t} \sum_{r \in Q} \gamma_{j,r} \sum_{p \in P_r} h'_{r,p} \beta_{r,p} \alpha_{r,p} \eta_{ij}, \quad (2.26)$$

where $\alpha_t = \alpha(h_t)$, $\alpha_{r,p} = \alpha(h_{r,p})$ (see (2.2)).

Similarly, still assuming that $i \notin Q$ we get

$$\mathcal{A}(g_t) = g_t + \sum_{r \in Q} \hat{g}_r \bar{\gamma}_{j,r} \eta_{ij}, \quad (2.27)$$

where $\hat{g}_r$ is obtained from g_t by replacing the vector e_j in the box (j, r) to ε_i ;

$$\bar{\gamma}_{j,r} = (-1)^{\bar{n}_{j,r}}, \quad (2.28)$$

where $\bar{n}_{j,r}$ is the number of odd base vectors that are placed after the g_t -box (j, r) when we move by the tableau from left to right and from the top down. Since the tableaux of h_t and g_t have the same filling within the small rectangle and the $(m+1)$ -th g_t -row of length n is filled with the odd base vectors $\varepsilon_1, \dots, \varepsilon_n$, we have

$$\bar{n}_{j,r} = n_{j,r} + n. \quad (2.29)$$

With (2.28), (2.29), (2.22) equality (2.27) takes the form

$$\mathcal{A}(g_t) = g_t + \sum_{r \in Q} \hat{g}_r \gamma_{j,r} (-1)^n \eta_{ij},$$

where $Q = Q_{g_t} = Q_{h_t} \setminus \{n+1\}$.

The tensors $\hat{g}_r$ are not canonical because the element ε_i is placed not in its “native” i -th column but in the r -th column ($r \neq i$).

To express $\hat{g}_r$ in terms of canonical tensors we use again the Jacobi identity. By analogy with (2.23), writing this identity for the i -th column and the box (j, r) we get

$$\hat{g}_r = -(q+1)\hat{g}_r + \sum_{p \in P_r} g_{r,p} \bar{\beta}_{r,p}, \quad (2.30)$$

where $g_{r,p}$ is obtained from $\hat{g}_r$ in the same way as $h_{r,p}$ is obtained from $\hat{h}_r$ (see above);

$$\bar{\beta}_{r,p} = (-1)^{\bar{m}_{r,p}},$$

where $\bar{m}_{r,p}$ is defined by analogy with $m_{r,p}$ (see above) but now for $\hat{g}_r$. Since the tableaux of $\hat{h}_r$ and $\hat{g}_r$ have the same filling within the small rectangle, we have $\bar{m}_{r,p} = m_{r,p}$. Hence,

$$\bar{\beta}_{r,p} = \beta_{r,p}.$$

From (2.30) it follows that

$$\hat{g}_r = \frac{1}{q+2} \sum_{p \in P_r} g_{r,p} \beta_{r,p}. \quad (2.31)$$

With (2.31) equality (2.27) takes the form

$$\mathcal{A}(g_t) = g_t + \frac{(-1)^n}{q+2} \sum_{r \in Q} \gamma_{j,r} \sum_{p \in P_r} g_{r,p} \beta_{r,p} \eta_{ij}. \quad (2.32)$$

We see that the canonical tensors $h_{r,p}$ and $g_{r,p}$ are corresponding in that these have the same filling of the small rectangle.

When (2.26) is compared with (2.32), it is apparent that to conclude the proof it remains to check the equality

$$\frac{\alpha_{r,p}}{\alpha_t} = \frac{(-1)^n(q+1)}{q+2}. \quad (2.33)$$

With (2.3) the last equality is equivalent to

$$\frac{(-1)^{n\rho(h_{r,p})} \kappa(h_{r,p}) \kappa(g_t)}{\kappa(g_{r,p}) (-1)^{n\rho(h_t)} \kappa(h_t)} = \frac{(-1)^n(q+1)}{q+2}. \quad (2.34)$$

First note that $\rho(h_{r,p}) - \rho(h_t) = 1$.

Further, for an arbitrary $z \in \{1, \dots, n\}$, $z \neq i$ the number of odd base vectors placed in the z -th column is the same for h_t and $h_{r,p}$ (also for g_t and $g_{r,p}$). The numbers of odd base vectors in the i -th column of h_t , $h_{r,p}$, g_t , $g_{r,p}$ are equal to q , $q + 1$, $q + 1$, $q + 2$ respectively. Then we have

$$\frac{\kappa(h_{r,p})\kappa(g_t)}{\kappa(g_{r,p})\kappa(h_t)} = \frac{(q+1)!(q+1)!}{(q+2)!q!} = \frac{q+1}{q+2}.$$

Hence (2.34) is proved.

Thus Lemma 2.2 is proved for the case when $i \notin Q$.

Now suppose $i \in Q$, that is, the h_t -box (j, i) is filled with e_j . Still we have (2.21). The tensors $\hat{h}_r$ are not canonical except when $r = i$.

The tensor $\hat{h}_{n+1}$ is not canonical because the element ε_i is in the box $(j, n+1)$, but the $(n+1)$ -th column must be filled with even base vectors only. To express $\hat{h}_{n+1}$ in terms of canonical tensors we use the Jacobi identity for the i -th column and the box $(j, n+1)$. In such a manner we get

$$\hat{h}_{n+1} = -q\hat{h}_{n+1} + \bar{h}\beta_{n+1,j}, \quad (2.35)$$

where as above q is the number of ε_i placed in the i -th h_t -column; $\bar{h}$ is obtained from $\hat{h}_{n+1}$ by the transposition of e_j and ε_i placed in the $\hat{h}_{n+1}$ -boxes (j, i) and $(j, n+1)$ (respectively).

Evidently, $\bar{h} = \hat{h}_i$. Also,

$$\beta_{n+1,j} = (-1)^{m_{n+1,j}},$$

where $m_{n+1,j}$ is the number of odd base vectors placed in the j -th $\hat{h}_{n+1}$ -row strictly between i -th and $(n+1)$ -th columns.

Now it follows from (2.35) that

$$\hat{h}_{n+1} = \frac{\beta_{n+1,j}}{q+1} \hat{h}_i. \quad (2.36)$$

Consider an arbitrary tensor $\hat{h}_r$, where $r \in Q \setminus \{i\}$. We claim that $j \in P_r$, where P_r is the set defined above. Indeed, by assumption the $\hat{h}_r$ -box (j, i) is filled with e_j , also ε_i is in the box (j, r) and e_j can be placed only in the j -th row in the tableau of $\hat{h}_r$.

By applying the Jacobi identity to the i -th column and the box (j, r) of the tableau of $\hat{h}_r$, we get

$$\hat{h}_r = -q\hat{h}_r + \sum_{p \in P_r \setminus \{j\}} h_{r,p}\beta_{r,p} + \hat{h}_i\beta_{r,j}, \quad (2.37)$$

where $h_{r,p}$, $\beta_{r,p}$ are defined above.

Recall that $\hat{h}_r$ is canonical iff $r = i$. By this reason (2.21) is conveniently rewritten in the form:

$$\mathcal{A}(h_t) = h_t + \sum_{r \in Q \setminus \{i\}} \hat{h}_r \gamma_{j,r} \eta_{ij} + \hat{h}_i \gamma_{j,i} \eta_{ij} + \hat{h}_{n+1} \gamma_{j,n+1} \eta_{ij}. \quad (2.38)$$

With (2.36) and (2.37) equality (2.38) takes the form:

$$\begin{aligned} \mathcal{A}(h_t) = h_t + \frac{1}{q+1} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \sum_{p \in P_r \setminus \{j\}} h_{r,p} \beta_{r,p} \eta_{ij} + \\ \sum_{r \in Q \setminus \{i\}} \frac{\gamma_{j,r}}{q+1} \hat{h}_i \beta_{r,j} \eta_{ij} + \hat{h}_i \gamma_{j,i} \eta_{ij} + \frac{\beta_{n+1,j}}{q+1} \hat{h}_i \gamma_{j,n+1} \eta_{ij}. \end{aligned} \quad (2.39)$$

Clearly (2.39) is equivalent to

$$\begin{aligned} \mathcal{A}(h_t) = h_t + \frac{1}{q+1} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \sum_{p \in P_r \setminus \{j\}} h_{r,p} \beta_{r,p} \eta_{ij} + \\ \hat{h}_i \left(\frac{1}{q+1} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \beta_{r,j} + \gamma_{j,i} + \frac{\beta_{n+1,j}}{q+1} \gamma_{j,n+1} \right) \eta_{ij}. \end{aligned} \quad (2.40)$$

Notice that $h_{r,p} \neq \hat{h}_i$, where $r \in Q \setminus \{i\}$, $p \in P_r \setminus \{j\}$.

We have

$$\gamma_{j,r} \beta_{r,j} = \gamma_{j,i}, \quad \gamma_{j,n+1} \beta_{n+1,j} = \gamma_{j,i}. \quad (2.41)$$

Then the coefficient at $\hat{h}_i$ in (2.40) is equal to

$$\frac{\gamma_{j,i}(l+q+2)}{q+1},$$

where $l = |Q|$. Therefore equality (2.40) takes the form

$$\mathcal{A}(h_t) = h_t + \frac{1}{q+1} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \sum_{p \in P \setminus \{j\}} h_{r,p} \beta_{r,p} \eta_{ij} + \frac{l+q+2}{q+1} \hat{h}_i \gamma_{j,i} \eta_{ij}.$$

Going to the “primed” tensors (see (2.3)) we obtain

$$\begin{aligned} \mathcal{A}(h'_t) = h'_t + \frac{1}{(q+1)\alpha_t} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \sum_{p \in P \setminus \{j\}} h'_{r,p} \beta_{r,p} \alpha_{r,p} \eta_{ij} + \\ \frac{l+q+2}{(q+1)\alpha_t} \hat{h}'_i \gamma_{j,i} \hat{\alpha}_i \eta_{ij}, \end{aligned} \quad (2.42)$$

where $\hat{\alpha}_i = \alpha(\hat{h}_i)$.

In its turn by analogy with (2.27) we have

$$\mathcal{A}(g_t) = g_t + \sum_{r \in Q \setminus \{i\}} \hat{g}_r \gamma_{j,r} \eta_{ij} (-1)^n + \hat{g}_i \gamma_{j,i} \eta_{ij} (-1)^n, \quad (2.43)$$

where $\hat{g}_r, \gamma_{j,r}, \gamma_{j,i}$ are defined as above for the case when $i \notin Q$. An important point is that the tensors $\hat{g}_r$ are not canonical for $r \in Q \setminus \{i\}$ and $\hat{g}_i$ is canonical.

To express an arbitrary tensor $\hat{g}_r$, where $r \in Q \setminus \{i\}$, in terms of canonical tensors we use the Jacobi identity for the i -th column and the box (j, r) . As it is mentioned above the condition $i \in Q$ implies $j \in P_r$. Then by analogy with (2.37), (2.30) we get

$$\hat{g}_r = -(q+1)\hat{g}_r + \sum_{p \in P_r \setminus \{j\}} g_{r,p} \beta_{r,p} + \hat{g}_i \beta_{r,j},$$

From the last equality it follows that

$$\hat{g}_r = \frac{1}{q+2} \sum_{p \in P_r \setminus \{j\}} g_{r,p} \beta_{r,p} + \frac{1}{q+2} \hat{g}_i \beta_{r,j}.$$

With the last expression for $\hat{g}_r$ equality (2.43) becomes:

$$\begin{aligned} \mathcal{A}(g_t) = g_t + \frac{(-1)^n}{q+2} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \sum_{p \in P \setminus \{j\}} g_{r,p} \beta_{r,p} \eta_{ij} + \\ \hat{g}_i (-1)^n \left(\frac{1}{q+2} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \beta_{r,j} + \gamma_{j,i} \right) \eta_{ij}. \end{aligned} \quad (2.44)$$

Using (2.41) we obtain

$$\frac{1}{q+2} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \beta_{r,j} + \gamma_{j,i} = \frac{\gamma_{j,i}(l+q+2)}{q+2}. \quad (2.45)$$

With (2.45) equality (2.44) takes the form

$$\begin{aligned} \mathcal{A}(g_t) = g_t + \frac{(-1)^n}{q+2} \sum_{r \in Q \setminus \{i\}} \gamma_{j,r} \sum_{p \in P \setminus \{j\}} g_{r,p} \beta_{r,p} \eta_{ij} + \\ \hat{g}_i (-1)^n \gamma_{j,i} \frac{l+q+2}{q+2} \eta_{ij}. \end{aligned} \quad (2.46)$$

Note that (2.33) is correct as above. Then comparison of (2.42) and (2.46) shows that to complete the proof of the lemma it suffices to check the equality

$$\frac{\hat{\alpha}_i}{\alpha_t} = \frac{(-1)^n(q+1)}{q+2}.$$

But the last one is correct for the same reason as (2.33) is.

Lemma 2.2 is completely proved.

Suppose $A_{h'}$ and A_g are as above.

Lemma 2.3. *If the matrix A of $\mathcal{A} \in GL V$ has the form (2.7), then*

$$A_{h'} = A_g.$$

For any matrix A of the form (2.7) we have $\text{Ber } A = 1$. Hence, according to Proposition 2.1, Lemma 2.3 is the next part of the proof of Theorem 2.1 (see (2.20)).

PROOF OF LEMMA 2.3. By assumption, there exist $i \in \{1, \dots, m\}$, $j \in \{1, \dots, n\}$ such that

$$\mathcal{A}(\varepsilon_j) = \varepsilon_j + e_i \xi_{ij},$$

where $\xi_{ij} \in G_1$, and for any $w \in \Omega \setminus \{\varepsilon_j\}$ we have $\mathcal{A}(w) = w$.

Let h_t be an element of Λ_h . By R denote the subset of $\{1, \dots, m\}$ such that $r \in R$ if and only if ε_j belongs to the box (r, j) of the h_t -tableau. Then we have

$$\mathcal{A}(h_t) = h_t + \sum_{r \in R} (-1)^{n_{r,j}} \tilde{h}_r \xi_{ij}, \quad (2.47)$$

where $\tilde{h}_r$ is obtained from h_t using the replacement of ε_j in the box (r, j) by e_i ; as above, $n_{r,j}$ is the number of odd base vectors that are placed from the box $(r, j+1)$ to the box $(m, n+1)$ when we move from left to right and from the top down by the rows of the h_t -tableau.

Suppose $i \in R$, that is, the h_t -box (i, j) is filled with ε_j . Since h_t is canonical, we see that the j -th h_t -column does not contain e_i .

Evidently, the tensor $\tilde{h}_i$ is canonical. We claim that it is the only canonical tensor among $\tilde{h}_r$, where $r \in R$. Indeed, consider the j -th $\tilde{h}_r$ -column, where $r \neq i$. Then e_i is placed not in its “native” i -th row but in the r -th row.

To reduce these tensors to a canonical form we use their skew-symmetry by column elements (see (1.4)). To be precise for a given $r \in R$ we transpose the vectors e_i and ε_j placed in the $\tilde{h}_r$ -boxes (i, j) and (r, j) respectively.

As a result of this transposition the additional factor $(-1)^{m_r+1}$ appears, where m_r is the number of odd base vectors placed strictly between the h_t -boxes (r, j) and (i, j) through the movement by the rows from left to right and from top to bottom.

Since

$$(-1)^{m_r+1} = (-1)^{n_{i,j}-n_{r,j}}$$

we see that all terms of the sum in the right-hand side of (2.47) are equal to $(-1)^{n_{i,j}} \tilde{h}_i \xi_{ij}$ and the number of these terms is $n_R = |R|$. Then equality (2.47) takes the form

$$\mathcal{A}(h_t) = h_t + (-1)^{n_{i,j}} n_R \tilde{h}_i \xi_{ij}.$$

Going to the “primed” tensors we obtain

$$\mathcal{A}(h'_t) = h'_t + (-1)^{n_{i,j}} n_R \tilde{h}'_i \frac{\alpha(\tilde{h}_i)}{\alpha(h_t)} \xi_{ij}. \quad (2.48)$$

If $i \notin R$ then the h_t -box (i, j) is filled with e_i . As a result all tensors $\tilde{h}_r$, where $r \in R$, are equal to zero because the element e_i enters twice in their j -th column but the tensors are skew-symmetric by elements of an arbitrary column. Thus for the case when $i \notin R$ we have

$$\mathcal{A}(h_t) = h_t. \quad (2.49)$$

Now let $g_t \in \Lambda_g$ be the tensor corresponding to $h_t \in \Lambda_h$ considered above, that is, the tableaux of h_t and g_t have the same filling within the small rectangle. By $\bar{R}$ denote the set of integers such that $r \in \bar{R}$ if and only if the g_t -box (r, j) is filled with ε_j . We claim that $\bar{R} = R \cup \{n+1\}$. In fact, since the tensor g_t is canonical, we see that the g_t -box $(m+1, j)$ is filled with ε_j and the tableaux of h_t and g_t are filled equally within the small rectangle.

Then we have

$$\mathcal{A}(g_t) = g_t + \sum_{r \in \bar{R}} (-1)^{\bar{n}_{r,j}} \tilde{g}_r \xi_{ij}, \quad (2.50)$$

where $\tilde{g}_r$ is obtained from g_t by the replacement of the element ε_j disposed in the g_t -box (r, j) to e_i ; $\bar{n}_{r,j}$ is the number of odd base vectors placed in the g_t -tableau strictly after the box (r, j) when we move by the rows from left to right and from top to bottom.

Evidently, we have $\bar{n}_{r,j} = n_{r,j} + n$ for $r \leq m$ and $\bar{n}_{m+1,j} = n - j$. Then separating the term for $r = m+1$ in right-hand side of (2.50) we obtain

$$\mathcal{A}(g_t) = g_t + \sum_{r \in R} (-1)^{n_{r,j}+n} \tilde{g}_r \xi_{ij} + (-1)^{n-j} \tilde{g}_{m+1} \xi_{ij}. \quad (2.51)$$

Suppose that $i \in \bar{R}$, that is, the element ε_j is placed in the g_t -box (i, j) and the element e_j does not enter in the j -th g_t -column.

The tensor $\tilde{g}_i$ is canonical and the rest tensors $\tilde{g}_r$, where $r \in \bar{R} \setminus \{i\}$, are not canonical because the element e_i is not placed in its “native” i -th row for these tensors.

To reduce a tensor $\tilde{g}_r$, where $r \in \bar{R} \setminus \{i\}$, to a canonical form we use skew-symmetry of the tensor by elements of the column (see (1.4)).

Then for any $r \in \bar{R} \setminus \{i\}$ we obtain the same tensor $\tilde{g}_i$ with the additional factor $(-1)^{\bar{m}_r+1}$, where $\bar{m}_r$ is the number of odd base vectors placed strictly between the boxes (r, j) and (i, j) (as above, we move from one box to another by the rows from left to right and from top to bottom). Since

$$(-1)^{\bar{m}_r+1} = (-1)^{n_{r,j}-n_{i,j}}$$

for $r \leq m$ and

$$(-1)^{\bar{m}_{m+1}+1} = (-1)^{n_{i,j}+j},$$

we see that all terms (distinct from g_t) in the right-hand side of (2.51) are equal to $(-1)^{n_{i,j}+n} \tilde{g}_i \xi_{ij}$ and the number of these terms is equal to $|\bar{R}| = n_R + 1$.

Then equality (2.50) takes the form

$$\mathcal{A}(g_t) = g_t + (-1)^{n_{i,j}+n}(n_R+1)\tilde{g}_i\xi_{ij}. \quad (2.52)$$

Comparison of (2.48) and (2.52) shows that to complete the proof of the lemma it suffices to check the equality

$$\frac{\alpha(\tilde{h}_i)}{\alpha(h_t)} = \frac{(-1)^n(n_R+1)}{n_R}. \quad (2.53)$$

By definition (see (2.2)) we have

$$\frac{\alpha(\tilde{h}_i)}{\alpha(h_t)} = \frac{(-1)^{n\rho(\tilde{h}_i)}\kappa(\tilde{h}_i)\kappa(g_t)}{\kappa(\tilde{g}_i)(-1)^{n\rho(h_t)}\kappa(h_t)}.$$

When passing from h_t to $\tilde{h}_i$ the number of odd base vectors within the small rectangle increases on 1, that is, $\rho(h_t) - \rho(h_i) = 1$.

The numbers of odd base vectors are the same for all respective columns of h_t and $\tilde{h}_i$ (g_t and $\tilde{g}_i$) except the j -th one. Also, the numbers of odd base vectors in the j -th column of h_t , $\tilde{h}_i$, g_t , $\tilde{g}_i$ are equal to n_R , $n_R - 1$, $n_R + 1$, n_R (respectively).

Hence we have

$$\frac{\kappa(\tilde{h}_i)\kappa(g_t)}{\kappa(\tilde{g}_i)\kappa(h_t)} = \frac{(n_R-1)!(n_R+1)!}{(n_R!)^2} = \frac{n_R+1}{n_R}.$$

and equality (2.53) is proved.

Suppose that $i \in \bar{R}$, that is, the box (i, j) is filled with e_i . By the same argument as above we get $\mathcal{A}(g_t) = g_t$. Also from (2.49) it follows that $\mathcal{A}(h'_t) = h'_t$. Thus the case when $i \in \bar{R}$ is trivial.

This concludes the proof of Lemma 2.3.

A further step in the proof of Theorem 2.1 is the following lemma.

Lemma 2.4. *If the matrix A of $\mathcal{A} \in GL V$ has the form (2.10), then*

$$A_{h'} = A_g.$$

PROOF OF LEMMA 2.4. By assumption, there exist unequal integers $s, t \in \{1, \dots, n\}$ such that

$$\mathcal{A}(\varepsilon_s) = \varepsilon_s + \varepsilon_t y,$$

where $y \in G_0$, and for any $w \in \Omega \setminus \{\varepsilon_s\}$ we have $\mathcal{A}(w) = w$.

Let h be an element of Λ_h . By F denote the set of integers such that an integer r belongs to F if and only if the h -box (r, s) is filled with ε_s . Suppose $F \neq \emptyset$. Denote

$$F_j = \{\nu \mid \nu \subseteq F, |\nu| = j\},$$

where $j = 0, 1, \dots, n_F$, $n_F = |F|$, that is, F_j is the set of all subsets of F with j elements. Then we have

$$\mathcal{A}(h) = h + \sum_{j=1}^{n_F} \sum_{\nu \in F_j} h_{j,\nu} y^j, \quad (2.54)$$

where $h_{j,\nu}$ is obtained from h using the replacement of ε_s placed in the h -boxes (i, s) , where $i \in \nu$, by ε_t .

For any $j > 0$ an arbitrary element $h_{j,\nu}$ is not canonical, because j elements of the form ε_t are placed not in their “native” t -th column but in the s -th column. For an arbitrary $h_{j,\nu}$ by $P(h_{j,\nu})$ denote the set of integers such that $p \in P(h_{j,\nu})$ if and only if e_p belongs to the t -th $h_{j,\nu}$ -column and e_p does not belong to the s -th $h_{j,\nu}$ -column. In other words, $p \in P(h_{j,\nu})$ iff the $h_{j,\nu}$ -box (p, t) is filled with e_p but the $h_{j,\nu}$ -box (p, s) is filled either with ε_s or with ε_t . Notice that for all $j > 0$, $\nu \in F_j$ the set $P(h_{j,\nu})$ is the same, that is, the composition of $P(h_{j,\nu})$ does not depend on j and ν . For this reason in what follows we write simply P instead of $P(h_{j,\nu})$.

Denote

$$P_j = \{\mu \mid \mu \subseteq P, |\mu| = j\},$$

where $j = 0, 1, \dots, n_F$, that is, P_j is the set of all subsets of P with j elements.

We claim that

$$h_{j,\nu} = \frac{1}{C_{q+j}^j} \sum_{\mu \in P_j} \delta_\mu \bar{h}_{j,\mu}, \quad (2.55)$$

where q is the number of odd base vectors (i.e., the number of ε_t) in the t -th column of h and $\bar{h}_{j,\mu}$ is obtained from $h_{j,\nu}$ as follows: for every $z \in \mu$ the elements e_z and ε_t placed in the $h_{j,\nu}$ -boxes (z, t) and (z, s) (respectively) change their places; $\delta_\mu = (-1)^{m_\mu}$, $m_\mu = \sum_{z \in \mu} m_z$, where m_z is the number of odd base vectors placed in the z -th $\bar{h}_{j,\mu}$ -row strictly between the boxes (z, t) and (z, s) . Note that we can assume that the $h_{j,\nu}$ -box (z, s) is filled with ε_t . Indeed, otherwise the box (z, s) is filled with ε_s and we use (1.4) to transpose ε_s and ε_t in the s -th column. Since the transposed elements are odd, we see that the additional sign does not appear.

Now let us prove identity (2.55). To express $h_{j,\nu}$ in terms of canonical tensors we apply the Jacobi identity step by step. In any step an element ε_t from the s -th column is moved to the t -th column. Also the application of the Jacobi identity in the next step brings the additional factor $1/(r+1)$, where r is the number of elements ε_t in the t -th column in the previous step. Thus at last the factor

$$\frac{1}{(q+1)(q+2)\cdots(q+j)}$$

arises. Further, there is one-to-one correspondence between the set of canonical tensors obtained in the last step and the set of all subsets of P with j elements. Also, any canonical tensor enters $j!$ times in the final expression for $h_{j,\nu}$, because an element e_z , $z \in P$, can come to the s -th column in any step of the algorithm and the number of these steps is equal to j .

Notice that the right-hand side of (2.55) depends on j and does not depend on ν . In other words, all tensors $h_{j,\nu}$, where $\nu \in F_j$, have the same representation in terms of the canonical tensors. Then with (2.55) equality (2.54) takes the form

$$\mathcal{A}(h) = h + \sum_{j=1}^{n_F} \frac{C_{n_F}^j}{C_{q+j}^j} \sum_{\mu \in P_j} \delta_\mu \bar{h}_{j,\mu} y^j$$

or in terms of the “primed” tensors we have

$$\mathcal{A}(h') = h' + \frac{1}{\alpha(h)} \sum_{j=1}^{n_F} \frac{C_{n_F}^j \alpha(h_{j,\mu})}{C_{q+j}^j} \sum_{\mu \in P_j} \delta_\mu \bar{h}'_{j,\mu} y^j, \quad (2.56)$$

where $h' = \frac{h}{\alpha(h)}$, $\bar{h}'_{j,\mu} = \frac{h_{j,\mu}}{\alpha(h_{j,\mu})}$.

Now let $g \in \Lambda_g$ be the tensor corresponding to h and $\mathcal{A} \in GL V$ as above. By $\bar{F}$ denote the set of integers such that $r \in \bar{F}$ if and only if the g -box (r, s) is filled with ε_s . Any column of g as compared with h contains the additional box filled with a proper odd base element (to be precise the g -box $(m+1, s)$ is filled with ε_s , where $s = 1, 2, \dots, n$). Hence we have

$$\bar{F} = F \cup \{m+1\}$$

and $n_{\bar{F}} = |\bar{F}| = n_F + 1$. Define

$$\bar{F}_j = \{\nu \mid \nu \subseteq \bar{F}, |\nu| = j\},$$

where $j = 0, 1, \dots, n_{\bar{F}}$, that is, $\bar{F}_j$ is the set of all subsets of $\bar{F}$ with j elements. Then we have

$$\mathcal{A}(g) = g + \sum_{j=1}^{n_F+1} \sum_{\nu \in \bar{F}_j} g_{j,\nu} y^j, \quad (2.57)$$

where $g_{j,\nu}$ are obtained from g just as $h_{j,\nu}$ are obtained from h (see above).

Since the tableaux of h and g have the same filling within the small rectangle and the $(m+1)$ -th g -row is filled with odd base vectors, we see that $P(h_{j,\nu}) = P(g_{j,\nu})$. Therefore we write P as above instead of $P(g_{j,\nu})$.

Let P_j be as above. We see that δ_μ , where $\mu \in P_j$, does not depend on either $h_{j,\nu}$ or $g_{j,\nu}$ is considered.

We have

$$g_{j,\nu} = \frac{1}{C_{q+j+1}^j} \sum_{\mu \in P_j} \delta_\mu \bar{g}_{j,\mu}, \quad (2.58)$$

where $\bar{g}_{j,\mu}$ are obtained from $g_{j,\nu}$ just as $\bar{h}_{j,\mu}$ are obtained from $h_{j,\nu}$. Indeed, by analogy with (2.55) equality (2.58) can be obtained and the only difference between these two cases is that $g_{j,\nu}$ has not q but $(q+1)$ elements ε_t in the t -th column. That is why the coefficient in (2.58) is equal not to $1/C_{q+j}^j$ (as in (2.55)) but to $1/C_{q+j+1}^j$.

With (2.58) equality (2.57) takes the form

$$\mathcal{A}(g) = g + \sum_{j=1}^{n_F} \frac{C_{n_F+1}^j}{C_{q+j+1}^j} \sum_{\mu \in P_j} \delta_\mu \bar{g}_{j,\mu} y^j + g_{\bar{F}} y^{n_F+1}, \quad (2.59)$$

where $g_{\bar{F}}$ is obtained from g using the replacement of *all* ε_s in the s -th g -column by ε_t . Let us show that

$$g_{\bar{F}} = 0. \quad (2.60)$$

Indeed, applying the Jacobi identity we represent $g_{\bar{F}}$ as a linear combination of tensors such that theirs s -th column is filled with even base vectors only. But the height of the s -th column is greater by 1 than the number m of even base vectors. Since any tensor of the form (1.3) is skew-symmetric by elements of any column we get (2.60).

Comparison of (2.56) and (2.59) shows that to complete the proof it suffices to check the equality

$$\frac{\alpha(\bar{h}_{j,\mu})}{\alpha(h)} = \frac{C_{q+j}^j C_{n_F+1}^j}{C_{n_F}^j C_{q+j+1}^j}$$

or, what is the same –

$$\frac{\alpha(\bar{h}_{j,\mu})}{\alpha(h)} = \frac{(n_F + 1)(q + 1)}{(n_F - j + 1)(q + j + 1)}. \quad (2.61)$$

We have:

	the numbers of odd base vectors	
	s -th column	t -th column
h	n_F	q
g	$n_F + 1$	$q + 1$
$h_{j,\nu}$	$n_F - j$	$q + j$
$\bar{g}_{j,\nu}$	$n_F - j + 1$	$q + j + 1$

Also note that the number of odd base vectors within the small rectangle is the same for the tableaux of h and $\bar{h}_{j,\mu}$, that is, we have $\rho(\bar{h}_{j,\mu}) = \rho(h)$. Hence we obtain

$$\begin{aligned} \frac{\alpha(\bar{h}_{j,\mu})}{\alpha(h)} &= \frac{\kappa(g)\kappa(\bar{h}_{j,\mu})}{\kappa(h)\kappa(\bar{g}_{j,\mu})} = \\ &= \frac{(n_F + 1)!(q + 1)!(n_F - j)!(q + j)!}{n_F!q!(n_F - j + 1)!(q + j + 1)!} \end{aligned}$$

and we see that (2.61) is proved.

This concludes the proof of Lemma 2.4.

Actually the following lemma is the last step in the proof of Theorem 2.1.

Lemma 2.5. *If the matrix A of $\mathcal{A} \in GL V$ has the form (2.9), then*

$$A_{h'} = A_g.$$

PROOF. By assumption, there exist integers $s, t \in \{1, 2, \dots, m\}$, $s \neq t$ and $x \in G_0$ such that

$$\mathcal{A}(e_s) = e_s + e_t x,$$

and for any $w \in \Omega \setminus \{e_s\}$ we have $\mathcal{A}(w) = w$.

Let h be an element of the set Λ_h . By H denote the set of integers such that $r \in H$ if and only if the h -boxes (s, r) and (t, r) are filled with e_s and ε_r (respectively). Evidently we have $(n+1) \notin H$. By H_j denote the set of all subsets of H with j elements, where $j = 1, 2, \dots, k_H$, $k_H = |H|$.

We have

$$\mathcal{A}(h) = h + \sum_{j=1}^{k_H} \sum_{\mu \in H_j} h_\mu x^j, \quad (2.62)$$

where h_μ is obtained from h when all elements e_s placed in the h -boxes (s, r) , where $r \in \mu$, are replaced by e_t .

An arbitrary tensor h_μ is not canonical in so far as the h_μ -boxes (s, r) , where $r \in \mu$, are filled with e_t but $t \neq s$.

To reduce a tensor h_μ , where $\mu \in H_j$, to a canonical form we use the skew-symmetry of this tensor by the elements of the columns. To be precise, for every $r \in \mu$ we transpose the vectors e_t and ε_r placed in the h_μ -boxes (s, r) and (t, r) respectively. The sign that appears as a result of this permutation we denote by σ_μ .

The canonical tensor obtained from h_μ is denoted by $\bar{h}_\mu$. Then equality (2.62) takes the form

$$\mathcal{A}(h) = h + \sum_{j=1}^{k_H} \sum_{\mu \in H_j} \sigma_\mu \bar{h}_\mu x^j,$$

and in terms of the “primed” tensors we have

$$\mathcal{A}(h') = h' + \frac{1}{\alpha(h)} \sum_{j=1}^{k_H} \sum_{\mu \in H_j} \sigma_\mu \bar{h}'_\mu \alpha(\bar{h}_\mu) x^j, \quad (2.63)$$

where $h' = h/\alpha(h)$, $\bar{h}'_\mu = \bar{h}_\mu/\alpha(h_\mu)$.

Note that for an arbitrary column the number of odd base vectors is the same for the tableaux of the tensors h , h_μ and $\bar{h}_\mu$. Hence we have $\alpha(\bar{h}_\mu) = \alpha(h)$, where $\mu \in H_j$, $j = 1, 2, \dots, k_H$. Thus (2.63) becomes

$$\mathcal{A}(h') = h' + \sum_{j=1}^{k_H} \sum_{\mu \in H_j} \sigma_\mu \bar{h}'_\mu x^j. \quad (2.64)$$

Now suppose g is the tensor corresponding to h , that is, g and h have the same filling within the small rectangle. Evidently the set H is the same for the tensors h and g . Then we get

$$\mathcal{A}(g) = g + \sum_{j=1}^{k_H} \sum_{\mu \in H_j} g_\mu x^j,$$

where g_μ is obtained from g much as h_μ is obtained from h . The tensors g_μ are reduced to a canonical form by the same process as h_μ are. Then we obtain

$$\mathcal{A}(g) = g + \sum_{j=1}^{k_H} \sum_{\mu \in H_j} \hat{\sigma}_\mu \bar{g}_\mu x^j, \quad (2.65)$$

where $\bar{g}_\mu$ are canonical tensors and we claim that $\hat{\sigma}_\mu = \sigma_\mu$. Indeed, g_μ -boxes touched by the permutation that reduces g_μ to a canonical form are placed strictly within the small rectangle.

Comparison of (2.64) and (2.65) shows that Lemma 2.5 is proved.

The case when the matrix A of $\mathcal{A}$ has the form (2.8) is trivial.

Thus we see that Theorem 2.1 is completely proved.

Let b_* be a formal element such that for any $\mathcal{A} \in GLV$ we have

$$\mathcal{A}(b_*) = (\text{Ber } A)^{-1} \cdot b_*,$$

where A is the matrix of $\mathcal{A}$.

Theorem 2.2. *The mapping*

$$\varphi_* : b_* \cdot h'_i \mapsto g_i$$

determines the isomorphism of GLV -modules $b_ \cdot W_{\lambda_h}$ and W_{λ_g} .*

PROOF. Follows immediately from Theorem 2.1.

3 An explicit construction of the one-dimensional representations of the general linear supergroup

By definition, put $\vartheta_i = e_i$ and $\vartheta_j = \varepsilon_{j-m}$, where $i = 1, 2, \dots, m$, $j = m+1, m+2, \dots, m+n$. The last notation gives the through enumeration of the set Ω .

By V^* denote the G -module dual to V .

Let l be a positive integer.

Suppose u_i^*, w_j are arbitrary homogeneous elements of V^* and V (respectively), where $i, j = 1, 2, \dots, l$. By definition, for the decomposable tensors $u_1^* \cdots u_l^* \in T_l(V^*)$, $w_1 \cdots w_l \in T_l(V)$ put

$$(u_1^* \cdots u_l^*, w_1 \cdots w_l) = (-1)^\chi (u_1^*, w_1) \cdots (u_l^*, w_l), \quad (3.1)$$

where χ is the number of pairs of odd elements that change their order when one pass from the sequence $u_1^*, \dots, u_l^*, w_1, \dots, w_l$ to the sequence $u_1^*, w_1, \dots, u_l^*, w_l$. Equality (3.1) determines the inclusion

$$T_l(V^*) \rightarrow (T_l(V))^*.$$

By definition, for arbitrary $u_1^*, \dots, u_l^* \in V^*$, $\sigma \in S_l$ put

$$(u_1^* \cdots u_l^*)\sigma = u_{\sigma(1)}^* \cdots u_{\sigma(l)}^*. \quad (3.2)$$

Thus $T_l(V^*)$ is the right S_l -module. With (1.2), (3.1), (3.2) we get

$$((u_1^* \cdots u_l^*)\sigma, w_1 \cdots w_l) = (u_1^* \cdots u_l^*, \sigma(w_1 \cdots w_l)), \quad (3.3)$$

where as above $u_i^* \in V^*$, $w_j \in V$.

By ϑ_i^* denote the elements of V^* dual to ϑ_j , that is,

$$(\vartheta_i^*, \vartheta_j) = \delta_{ij},$$

where $i, j = 1, 2, \dots, m+n$, δ_{ij} is the Kronecker delta. Clearly, $e_i^* = \vartheta_i^*$, $\varepsilon_j^* = \vartheta_{j+m}^*$, where $i = 1, \dots, m$, $j = 1, \dots, n$. By analogy with the notation used above, $\Omega_0^* = \{e_1^*, \dots, e_m^*\}$, $\Omega_1^* = \{\varepsilon_1^*, \dots, \varepsilon_n^*\}$, $\Omega^* = \Omega_0^* \cup \Omega_1^*$.

We say that $f \in T_l(V^*)$ is an Ω^* -tensor if there exist $u_1^*, \dots, u_l^* \in \Omega^*$ and a Young tableau T such that

$$f = (u_1^* \cdots u_l^*)e_T.$$

Let Λ_g be as above (see §1), g_i elements of Λ_g in some ordering of the last one, where $i = 1, 2, \dots, k_\Lambda$; $k_\Lambda = 2^{mn}$. For an arbitrary $i \in \{1, \dots, k_\Lambda\}$ by g_i^* denote the Ω^* -tensor such that g_i^* -tableau is obtained from g_i -tableau by the formal change $\vartheta_j \mapsto \vartheta_j^*$, where $j = 1, 2, \dots, (m+n)$.

By k_i denote the cardinality of the automorphism group of the g_i -tableau and by q_i denote the number of odd base elements in the g_i -tableau.

Lemma 3.1. *For arbitrary $i, j \in \{1, 2, \dots, k_\Lambda\}$ the following equality holds*

$$(g_i^*, g_j) = \delta_{ij} \zeta_i, \quad (3.4)$$

where

$$\zeta_i = k_i \mu_{\lambda_g} (-1)^{(q_i^2 - q_i)/2},$$

$$\mu_{\lambda_g} = \frac{\prod_{t=1}^{m+1} (m+n+1-t)!}{\prod_{t=1}^{m+1} (m+1-t)!}. \quad (3.5)$$

PROOF. By definition, there exist $u_1^*, \dots, u_l^* \in \Omega^*$, $w_1, \dots, w_l \in \Omega$ such that

$$g_i^* = (u_1^* \cdots u_l^*) e_T, \quad g_j = e_T(w_1 \cdots w_l),$$

where T is the Young tableau with the diagram $\lambda_g = (\underbrace{n, \dots, n}_{m+1})$, $l = (m+1)n$.

With (3.3) we get

$$(g_i^*, g_j) = ((u_1^* \cdots u_l^*) e_T, e_T(w_1 \cdots w_l)) = (u_1^* \cdots u_l^*, e_T^2(w_1 \cdots w_l)).$$

For an arbitrary Young diagram λ the following identity holds

$$e_{T_\lambda}^2 = \mu_\lambda e_{T_\lambda},$$

where T_λ is a Young tableau with the diagram λ , μ_λ is a non-zero integer. In particular, μ_{λ_g} is given by (3.5) (see [6]). By assumption, $u_1^*, \dots, u_l^*$, $w_1, \dots, w_l$ fill the diagram λ_g in a canonical way. Then to conclude the proof it suffices to use Lemma 1.1 from [2].

Thus Lemma 3.1 is proved.

An arbitrary linear transformation $\mathcal{A} \in GLV$ determines, as it usually does, the corresponding linear transformation of V^* by the equalities

$$(\mathcal{A}(\vartheta_i^*), \mathcal{A}(\vartheta_j)) = (\vartheta_i^*, \vartheta_j) = \delta_{ij} \quad (3.6)$$

(this transformation of V^* we still denote by $\mathcal{A}$).

Suppose A is the matrix of $\mathcal{A}$, that is, we have

$$(\mathcal{A}(\vartheta_1), \dots, \mathcal{A}(\vartheta_{m+n})) = (\vartheta_1, \dots, \vartheta_{m+n}) \cdot A. \quad (3.7)$$

Then with (3.6) we obtain

$$\begin{pmatrix} \mathcal{A}(\vartheta_1^*) \\ \mathcal{A}(\vartheta_2^*) \\ \dots \\ \mathcal{A}(\vartheta_{m+n}^*) \end{pmatrix} = A^{-1} \cdot \begin{pmatrix} \vartheta_1^* \\ \vartheta_2^* \\ \dots \\ \vartheta_{m+n}^* \end{pmatrix}. \quad (3.8)$$

Denote

$$g_i^{*'} = \frac{1}{(g_i^*, g_i)} g_i^*.$$

From (3.4) it follows that

$$(g_i^{*'}, g_j) = \delta_{ij}.$$

Recall that $\mathcal{A}$ is even. Then it follows from (3.1) and (3.6) that

$$(\mathcal{A}(g_i^{*'}), \mathcal{A}(g_j)) = (g_i^{*'}, g_j) = \delta_{ij}.$$

In other words, the action of $\mathcal{A}$ on $\langle g_i \rangle$ and $\langle g_i^{*'} \rangle$ is the particular case of changing base in a space and the dual one (see (3.7), (3.8)).

Consequently we have

$$\begin{aligned} \sum_{i=1}^{k_\Lambda} \mathcal{A}(g_i) \mathcal{A}(g_i^{*'}) &= (\mathcal{A}(g_1), \dots, \mathcal{A}(g_{k_\Lambda})) \begin{pmatrix} \mathcal{A}(g_1^{*'}) \\ \dots \\ \mathcal{A}(g_{k_\Lambda}^{*'}) \end{pmatrix} = \\ &= (g_1, \dots, g_{k_\Lambda}) \tilde{A} \tilde{A}^{-1} \begin{pmatrix} g_1^{*'} \\ \dots \\ g_{k_\Lambda}^{*'} \end{pmatrix} = (g_1, \dots, g_{k_\Lambda}) \begin{pmatrix} g_1^{*'} \\ \dots \\ g_{k_\Lambda}^{*'} \end{pmatrix} = \sum_{i=1}^{k_\Lambda} g_i g_i^{*'}. \end{aligned}$$

This completes the proof of the following result:

Proposition 3.1. *The action of an arbitrary $\mathcal{A} \in GLV$ on the tensor*

$$\sum_{i=1}^{k_\Lambda} g_i g_i^{*'}$$

is identical.

Theorem 3.1. *The tensor*

$$\tilde{b} = \sum_{i=1}^{k_\Lambda} h'_i g_i^{*'}$$

generates the one-dimensional GLV -module such that

$$\mathcal{A}(\tilde{b}) = \text{Ber } A \cdot \tilde{b},$$

where A is a matrix of $\mathcal{A} \in GLV$.

PROOF. Follows immediately from Theorem 2.1 and Proposition 3.1.

Consider some examples.

Example 3.1. Suppose $n = 0$. In this case one can say that the small rectangle degenerates into the segment of height m . Then $\tilde{b} = \gamma h_1$, where h_1 is the tensor skew-symmetric in $e_1, \dots, e_m$; γ is an invertible element of G_0 . Also, for $n = 0$ we have $\text{Ber } A = \det A$. Thus we arrive at the basic classic result.

Example 3.2. Let the dimension of V be $1|1$; $V_0 = \langle e \rangle$, $V_1 = \langle \varepsilon \rangle$. By $\mathcal{T}_{h_i}$ ($\mathcal{T}_{g_i}$) denote the h_i -tableau (g_i -tableau, respectively). Then we have

$$\mathcal{T}_{h_1} = \begin{bmatrix} e & e \end{bmatrix}, \quad \mathcal{T}_{h_2} = \begin{bmatrix} \varepsilon & e \end{bmatrix},$$

$$h_1 = (\mathbf{e} + (12))ee = 2ee,$$

$$h_2 = (\mathbf{e} + (12))\varepsilon e = \varepsilon e + e\varepsilon,$$

where $\mathbf{e}$ is the unit of S_2 . With (2.2) and (2.3) we get

$$h'_1 = 2ee, \quad h'_2 = -2(\varepsilon e + e\varepsilon). \quad (3.9)$$

Further, we have

$$\mathcal{T}_{g_1} = \begin{bmatrix} e \\ \varepsilon \end{bmatrix}, \quad \mathcal{T}_{g_2} = \begin{bmatrix} \varepsilon \\ \varepsilon \end{bmatrix},$$

$q_1 = k_1 = 1$, $q_2 = k_2 = 2$, $\mu_{\lambda_g} = 2$. Hence, $\zeta_1 = 2$, $\zeta_2 = -4$. Therefore,

$$g_1^{*'} = \frac{1}{2}(e^*\varepsilon^*(\mathbf{e} - (12))) = \frac{1}{2}(e^*\varepsilon^* - \varepsilon^*e^*), \quad (3.10)$$

$$g_2^{*'} = -\frac{1}{4}(\varepsilon^*\varepsilon^*(\mathbf{e} - (12))) = -\frac{1}{2}\varepsilon^*\varepsilon^*. \quad (3.11)$$

Now with (3.9), (3.10), (3.11) we obtain

$$\tilde{b} = \sum_{i=1}^2 h'_i g_i^{*'} = eee^*\varepsilon^* - ee\varepsilon^*e^* + e\varepsilon\varepsilon^*\varepsilon^* + \varepsilon e\varepsilon^*\varepsilon^*.$$

Let $\Lambda_h = \{h_i\}$ be as above (see §1). For an arbitrary $i \in \{1, 2, \dots, k_\Lambda\}$ by h_i^* denote the Ω -tensor such that the h_i^* -tableau is obtained from the h_i -tableau by the formal change $\vartheta_j \mapsto \vartheta_j^*$, where $j = 1, 2, \dots, (m+n)$.

By l_i denote the cardinality of the automorphism group of the h_i -tableau and by p_i denote the number of odd elements of the h_i -tableau.

Then by analogy with Lemma 3.1 we obtain:

Lemma 3.2. *For arbitrary $i, j \in \{1, 2, \dots, k_\Lambda\}$ the following equality holds*

$$(h_i^*, h_j) = \delta_{ij} \zeta'_i,$$

where

$$\begin{aligned} \zeta'_i &= l_i \mu_{\lambda_h} (-1)^{(p_i^2 - p_i)/2}, \\ \mu_{\lambda_h} &= \frac{\prod_{t=1}^m (m+n+1-t)!}{\prod_{t=1}^m (m-t)!}. \end{aligned}$$

By definition, put

$$h_i^{*'} = \frac{\alpha(h_i)}{(h_i^*, h_i)} h_i^*.$$

With (2.3) we have

$$(h_i^{*'}, h_j') = \delta_{ij}.$$

By analogy with Proposition 3.1 and Theorem 3.1 we arrive at the following results:

Proposition 3.2. *The action of an arbitrary $\mathcal{A} \in GL V$ on the tensor*

$$\sum_{i=1}^{k_\Lambda} h_i' h_i^{*'}.$$

is identical.

Theorem 3.2. *The tensor*

$$\tilde{b}_* = \sum_{i=1}^{k_\Lambda} g_i h_i^{*'}.$$

generates the one-dimensional GLV -module such that

$$\mathcal{A}(\tilde{b}_*) = (\text{Ber } A)^{-1} \cdot \tilde{b}_*,$$

where A is a matrix of $\mathcal{A} \in GLV$.

Example 3.3. Let the dimension of V be $1|1$; $V_0 = \langle e \rangle$, $V_1 = \langle \varepsilon \rangle$. Then we have

$$h_1^* = e^* e^* (\mathbf{e} + (12)) = 2e^* e^*,$$

$$h_2^* = \varepsilon^* e^* (\mathbf{e} + (12)) = \varepsilon^* e^* + e^* \varepsilon^*$$

(see the h_i -tableaux $\mathcal{T}_{h_i}$ in Example 3.2). Also, $l_1 = 2$, $p_1 = 0$, $l_2 = p_2 = 1$, $\mu_{\lambda_h} = 2$. Hence, $\zeta'_1 = 4$, $\zeta'_2 = 2$.

With (2.3) we get $\alpha(h_1) = 1$, $\alpha(h_2) = -\frac{1}{2}$. Thus we obtain

$$h_1^{*'} = \frac{1}{2} e^* e^*, \quad h_2^{*'} = -\frac{1}{4} (\varepsilon^* e^* + e^* \varepsilon^*).$$

Further,

$$g_1 = (\mathbf{e} - (12))e\varepsilon = e\varepsilon - \varepsilon e,$$

$$g_2 = (\mathbf{e} - (12))\varepsilon\varepsilon = 2\varepsilon\varepsilon$$

(see the g_i -tableaux $\mathcal{T}_{g_i}$ in Example 3.2). At last, we obtain

$$\tilde{b}_* = \sum_{i=1}^2 g_i h_i^{*'} = \frac{1}{2} (e\varepsilon e^* e^* - \varepsilon e e^* e^* - \varepsilon \varepsilon \varepsilon^* e^* - \varepsilon \varepsilon e^* \varepsilon^*).$$

References

- [1] H.M. Khudaverdian and Th.Th. Voronov, Berezinians, exterior powers and recurrent sequences, Lett. Math. Phys., Berezin's memorial volume, November, 2005.
- [2] I. Kantor, I. Trishin, On a Concept of Determinant in the Supercase, Comm. in Algebra, 22 (10), (1994), 3679-3739.

- [3] I.M. Trishin, On reduction of elements of the full matrix superalgebra to a block diagonal form by conjugation, *Linear Algebra Appl.* 357 (2002), 59-82.
- [4] F.A. Berezin, *Introduction to Superanalysis*, Reidel, Dordrecht, 1987.
- [5] I.L. Kantor, I.M. Trishin, Identical relations for the forms with the Young symmetry, *Geometriae Dedicata* 45: 307-321, 1993.
- [6] Weyl H., *The Classical Groups: Their Invariants and Representations*, Second edition, Princeton University Press, N.Y., 1997 (fifteenth printing).